

# Interstellar extinction and polarization — A spheroidal dust grain approach perspective

H.K. Das<sup>1\*</sup>, N.V. Voshchinnikov<sup>2</sup> and V.B. Il'in<sup>2,3</sup>

<sup>1</sup>*IUCAA, Post Bag 4, Ganeshkhind, Pune 411 007, India*

<sup>2</sup>*Sobolev Astronomical Institute, St. Petersburg University, St. Petersburg 198504, Russia*

<sup>3</sup>*Main (Pulkovo) Astronomical Observatory, St. Petersburg 196140, Russia*

Received 2009 ...

## ABSTRACT

We extend and investigate the spheroidal model of interstellar dust grains used to simultaneously interpret the observed interstellar extinction and polarization curves. We compare our model with similar models recently suggested by other authors, study its properties and apply it to fit the normalized extinction  $A(\lambda)/A_V$  and the polarizing efficiency  $P(\lambda)/A(\lambda)$  measured in the near IR to far UV region for several stars seen through one large cloud. We conclude that the model parameter  $\Omega$  being the angle between the line of sight and the magnetic field direction can be more or less reliably determined from comparison of the theory and observations. This opens a way to study the spatial structure of interstellar magnetic fields by using multi-wavelength photometric and polarimetric observations.

**Key words:** ISM: clouds – polarization – dust, extinction.

## 1 INTRODUCTION

Modelling of the wavelength dependencies of interstellar extinction  $A(\lambda)$  and polarization  $P(\lambda)$  along with analysis of infrared bands is the main source of information about the properties of interstellar grains (Draine 2003; Whittet 2003; Voshchinnikov 2004).

Extinction data were often used to constrain the size distribution of carbonaceous and silicate bare grains (e.g. Mathis, Rumpl & Nordsieck 1977; Zubko, Krelowski & Wegner 1996; Weingartner & Draine 2001). These data were also modeled with composite particles (Vaidya et al. 2001; Voshchinnikov et al. 2006; Iati et al. 2008). In all these studies the grains were assumed to be spherical, i.e. their shape was not involved.

To get information about the grain shapes as well as the ambient magnetic fields one needs to consider interstellar polarization. Interpretation of its wavelength dependence includes calculations of the extinction cross sections of rotating partially aligned non-spherical particles. In early studies (e.g. Hong & Greenberg 1980; Voshchinnikov 1989) an unphysical model of infinitely long cylinders was applied<sup>1</sup>. More realistic is the spheroidal model of

grains with the shape of these axisymmetric particles being characterized by the only parameter — the ratio of the major to minor semi-axis  $a/b$ . The optical properties of spheroids have long been of interest (e.g., Martin 1978; Rogers & Martin 1979; Onaka 1980; Draine & Lee 1984; Voshchinnikov & Farafonov 1993; Kim & Martin 1995). The growth of such grains in the interstellar medium (ISM) has been discussed by Stark, Potts & Diver (2006), their absorption and scattering properties (in the Rayleigh limit) by Min et al. (2006) and mineralogy by Li, Zhao & Li (2007).

Over the past decade spheroids have found wide applications in studies of optics of cometary dust — Das & Sen (2006), Moreno et al. (2007), circumstellar dust — Wolf et al. (2002), interplanetary dust — Lasue et al. (2007). Greenberg & Li (1996) used prolate spheroids and Lee & Draine (1985), Hildebrand & Dragovan (1995) oblate ones to interpret the silicate and ice polarization features of the BN object. Draine & Allaf-Akbari (2006) predicted halos due to X-ray scattering by oblate particles, detection of which would serve as a test for dust models.

The spheroidal model of cosmic dust grains is particularly promising for interpretation of the interstellar polarization and extinction data. A detailed review of early works on the subject can be found in Voshchinnikov (2004), so we consider recent applications of the model. Gupta et al. (2005) well reproduced the average Galactic extinction curve in the region  $0.3 - 9.5 \mu\text{m}^{-1}$  with silicate and graphite oblate spheroidal ( $a/b = 1.33$ ) particles rotating in a plane

\* E-mail: hkdas@iucaa.ernet.in (HKD); nvv@astro.spbu.ru (NVV); ilin55@yandex.ru (VBI)

<sup>1</sup> Li & Greenberg (1997) also considered finite cylinders for modelling interstellar extinction and polarization.

and having the size distribution  $n(r_V) \propto r_V^{-3.5}$  with  $r_V = 0.005 - 0.25 \mu\text{m}$  where  $r_V$  is the radius of a volume equivalent sphere. They found that the shape of grains did not affect the abundance problem, but essentially changed the polarization efficiency factors at different particle orientation angles. It should be noted that interpretation of the extinction data alone is usually made within the spherical model of dust grains (see for details the recent review of Draine 2009). Draine & Allaf-Akbari (2006) and Draine & Fraisse (2009) fitted both the average Galactic extinction and polarization curves in the range  $0.4 - 9.5 \mu\text{m}^{-1}$  applying picket-fence oriented silicate and graphite oblate spheroids with the aspect ratio in the interval  $a/b = 1 - 2$  with specific size distributions. The grain alignment degree was size dependent and showed a sharp increase at  $r_V \sim 0.1 \mu\text{m}$ . Voshchinnikov & Das (2008; hereafter VD08) studied the wavelength dependence of the ratio of the linear polarization degree to extinction (so called polarizing efficiency) from the ultraviolet to near-infrared. They found that the wavelength dependence of  $P(\lambda)/A(\lambda)$  was mainly determined by the particle composition and size whereas the absolute values of this ratio depended on the particle shape, degree and direction of alignment.

In this paper we investigate further advantages of the spheroidal approach to interstellar grains. In contrast to other works we calculate the optical properties of spheroids of all sizes exactly even at far UV wavelengths, consider ensembles of particles with various degree and direction of alignment, compare different approaches to extinction and polarization curve modelling, treat the observational data in a new way (the wavelength dependence of the polarizing efficiency is fitted instead of the polarization curve), apply the theory to observational data for concrete stars, and discuss a possibility to determine the direction of interstellar magnetic fields from the data. In Section 2 we describe details of our model and compare it with the models used earlier. In Section 3 the results obtained from our model are compared with observations of seven stars for which the line of sight intersects one main cloud, keeping in view that “mean extinction curves as well as mean total-to-selective extinction ratios are possibly averaged over several interstellar clouds of possibly different physical parameters” (Krelowski 1989). Section 4 contains our conclusions.

## 2 THEORY

Our modelling is based on the simple standard assumptions concerning dust grain materials, size distribution, shape and alignment. Homogeneous spheroids considered are characterized by their type (prolate or oblate), the aspect ratio  $a/b$  and a size parameter. As the latter we choose the radius  $r_V$  of a sphere whose volume is equal to that of a non-spherical particle, i.e.  $r_V^3 = ab^2$  for prolate spheroids and  $r_V^3 = a^2b$  for oblate ones.

### 2.1 Dust grain materials

Interstellar dust mainly consists of five heavy elements: C, O, Mg, Si and Fe, which are locked in carbonaceous and silicate particles (Jones 2000; Draine 2009). We consider sub-micron

grains from amorphous carbon (AC1) and amorphous silicate (astronomical silicate, astrosil). In order to reproduce the 2175Å absorption feature small graphite spheres with radius  $r_{\text{gra}} = 0.02 \mu\text{m}$  are also involved<sup>2</sup>. The optical constants of the materials are taken from Rouleau & Martin (1991) and Laor & Draine (1993).

### 2.2 Size distribution

Interpretation of the interstellar extinction is usually aimed at reconstruction of the dust grain size distribution consistent with the cosmic abundances and some other constraints (see discussion in Voshchinnikov 2004).

We choose a power-law size distribution

$$n(r_V) \propto r_V^{-q}, \quad (1)$$

which was derived by Mathis, Rumpl & Nordsieck (1977) (see also Draine & Lee 1984) from minimization of  $\chi^2$ -statistic. The distribution has three parameters: the lower ( $r_{V,\text{min}}$ ) and upper ( $r_{V,\text{max}}$ ) limits and the power index  $q$ . It was widely used in modelling of interstellar extinction and radiative transfer in dusty objects. The average Galactic extinction curve can be reproduced with the so-called standard MRN mixture — an ensemble of carbonaceous (graphite) and silicate spheres (in nearly equal proportions) with the parameters:  $q = 3.5$ ,  $r_{V,\text{min}} \approx 0.005 \mu\text{m}$  and  $r_{V,\text{max}} \approx 0.25 \mu\text{m}$ .

### 2.3 Cross section averaging

Let us consider non-polarized light passing through a cloud that contains three populations of grains: rotating spheroids from amorphous silicate (Si), amorphous carbon (C) and graphite (gra) spheres. For a line of sight, extinction (in stellar magnitudes) and linear polarization (in percentage) produced by the cloud can be written as

$$A(\lambda) = 1.086 N_d \langle C_{\text{ext}} \rangle_\lambda, \quad P(\lambda) = N_d \langle C_{\text{pol}} \rangle_\lambda 100\%. \quad (2)$$

Here,  $N_d = N_C + N_{\text{Si}} + N_{\text{gra}}$  is the total dust grain column density,  $\langle C_{\text{ext}} \rangle_\lambda$  and  $\langle C_{\text{pol}} \rangle_\lambda$  are the extinction and polarization cross sections, respectively, averaged over the grain populations, i.e.

$$\langle C_{\text{ext}} \rangle_\lambda = \mathcal{K}_C \overline{C}_{\text{ext},C}(\lambda) + \mathcal{K}_{\text{Si}} \overline{C}_{\text{ext,Si}}(\lambda) + \mathcal{K}_{\text{gra}} C_{\text{ext,gra}}(\lambda), \quad (3)$$

$$\langle C_{\text{pol}} \rangle_\lambda = \mathcal{K}_C \overline{C}_{\text{pol},C}(\lambda) + \mathcal{K}_{\text{Si}} \overline{C}_{\text{pol,Si}}(\lambda), \quad (4)$$


where  $C_{\text{ext,gra}}(\lambda) = \pi r_{\text{gra}}^2 Q_{\text{ext,gra}}(\lambda)$  and  $\mathcal{K}_i = N_i/N_d$  ( $i \equiv C, \text{Si}$ ) is the relative column density of the  $i$ -th population. It is evident that

$$\mathcal{K}_C + \mathcal{K}_{\text{Si}} + \mathcal{K}_{\text{gra}} = 1.$$

The values of  $\overline{C}_{\text{ext,pol},i}$  are obtained by averaging of the cross sections over the size distribution and grain orientations (Hong & Greenberg 1980; Voshchinnikov 1989)

$$\begin{aligned} \overline{C}_{\text{ext},i}(\lambda) &= \left(\frac{2}{\pi}\right)^2 \int_{r_{V,\text{min},i}}^{r_{V,\text{max},i}} \int_0^{\pi/2} \int_0^{\pi/2} \int_0^{\pi/2} \pi r_{V,i}^2 Q_{\text{ext},i}(\lambda) \\ &\times n_i(r_V) f(\xi, \beta) d\varphi d\omega d\beta dr_V, \end{aligned} \quad (5)$$

<sup>2</sup> Note that graphite is not the only material considered as the carrier of the 2175Å absorption feature (see, e.g., Li et al. 2008)



**Figure 1.** Geometrical configuration of a spinning and wobbling prolate spheroidal grain. The major (symmetry) axis of the particle  $O_1O_2$  is situated in the spinning plane  $NO_1O_2$  which is perpendicular to the angular momentum  $J$ . The direction of light propagation  $k$  is parallel to the  $Z$ -axis and makes the angle  $\alpha$  with the particle symmetry axis.

$$\begin{aligned} \overline{C}_{\text{pol},i}(\lambda) &= \frac{2}{\pi^2} \int_{r_{V,\min,i}}^{r_{V,\max,i}} \int_0^{\pi/2} \int_0^\pi \int_0^{\pi/2} \pi r_{V,i}^2 Q_{\text{pol},i}(\lambda) \\ &\times n_i(r_V) f(\xi, \beta) \cos 2\psi d\varphi d\omega d\beta dr_V. \end{aligned} \quad (6)$$

Here,  $\beta$  is the precession-cone angle for the angular momentum  $J$  which spins around the direction of the magnetic field  $B$ ,  $\varphi$  the spin angle,  $\omega$  the precession angle (see Fig. 1). The quantities  $Q_{\text{ext}} = (Q_{\text{ext}}^{\text{TM}} + Q_{\text{ext}}^{\text{TE}})/2$  and  $Q_{\text{pol}} = (Q_{\text{ext}}^{\text{TM}} - Q_{\text{ext}}^{\text{TE}})/2$  are the extinction and polarization efficiency factors for non-polarized incident radiation (Voshchinnikov & Farafonov 1993),

$$\cos 2\psi = \begin{cases} \cos 2\Phi \left[ \frac{2 \cos^2 \varphi \cos^2 \theta}{1 - \cos^2 \varphi \sin^2 \theta} - 1 \right] & , \text{ prolate} \\ \cos 2\Phi & , \text{ oblate,} \end{cases} \quad (7)$$

and

$$\cos \theta = \cos \Omega \cos \beta + \sin \Omega \sin \beta \cos \omega, \quad (8)$$

$$\cos 2\Phi = \left[ \frac{2 \sin^2 \beta \sin^2 \omega}{\sin^2 \theta} - 1 \right], \quad (9)$$

$$\cos \alpha = \begin{cases} \frac{\sin \theta \cos \varphi}{\sqrt{1 - \sin^2 \theta \cos^2 \varphi}} & , \text{ prolate} \\ \sqrt{1 - \sin^2 \theta \cos^2 \varphi} & , \text{ oblate.} \end{cases} \quad (10)$$

Here,  $\Omega$  is the angle between the line of sight and the magnetic field direction ( $0^\circ \leq \Omega \leq 90^\circ$ ). The value  $\Omega = 90^\circ$  corresponds to the case when the particle rotation plane contains the light propagation vector  $k$ , which gives the maximum degree of linear polarization. For  $\Omega = 0^\circ$ , the light falls perpendicular to the particle rotation plane and from symmetry reasons the net degree of polarization produced is zero.

The efficiency factors were calculated by using a solution to the light scattering problem for a homogeneous

spheroid obtained by the method of separation of variables in spheroidal coordinates (Voshchinnikov & Farafonov 1993) and a modern approach to computations of spheroidal wave functions (Voshchinnikov & Farafonov 2003). In contrast to all works where the spheroidal model was recently applied (Gupta et al. 2005; Draine 2009; Draine & Fraise 2009) we computed the optical properties of the particles of all sizes exactly even in the far UV spectral region.

## 2.4 Grain alignment

Interstellar non-spherical grains should be partially aligned. The “picket fence” or perfect rotating alignment often used in modelling of polarization (e.g. Kim & Martin 1995; Draine & Fraise 2009) is a crude approximation giving a too high polarizing efficiency in comparison with the empirical upper limit (e.g., Greenberg 1978)

$$P_{\text{max}}/A_V \lesssim 3\%/\text{mag}.$$

Although the theory of grain alignment is actively discussed (Lazarian 2007) it still has little practical significance. Therefore, we consider the classical alignment mechanism — so-called imperfect Davis–Greenstein (IDG) orientation (Davis & Greenstein 1951). The IDG mechanism is described by the function  $f(\xi, \beta)$  depending on the alignment parameter  $\xi$  and the precession angle  $\beta$  (Hong & Greenberg 1980; Voshchinnikov 1989)

$$f(\xi, \beta) = \frac{\xi \sin \beta}{(\xi^2 \cos^2 \beta + \sin^2 \beta)^{3/2}}. \quad (11)$$

The parameter  $\xi$  depends on the particle size  $r_V$ , the imaginary part of the grain magnetic susceptibility  $\chi'' = \kappa \omega_d / T_d$ , where  $\omega_d$  is the angular velocity of a grain and  $\kappa = 2.5 \times 10^{-12}$  (Davis & Greenstein 1951), gas density  $n_H$ , the strength of magnetic field  $B$ , and temperatures of dust  $T_d$  and gas  $T_g$ . By introducing a parameter

$$\delta_0 = 8.23 \times 10^{23} \frac{\kappa B^2}{n_H T_g^{1/2} T_d}, \quad \mu\text{m}, \quad (12)$$

one can get

$$\xi^2 = \frac{r_V + \delta_0 (T_d/T_g)}{r_V + \delta_0}. \quad (13)$$

We assume that carbon and silicate grains have similar alignment but different size distributions. Thus, our model has 10 main parameters:  $r_{V,\min}$ ,  $r_{V,\max}$  and  $q$  for carbon and silicate grains, relative density for carbon ( $\mathcal{K}_C$ ) and silicate ( $\mathcal{K}_{Si}$ ) grains, degree ( $\delta_0$ ) and direction ( $\Omega$ ) of grain alignment.

## 2.5 Comparison with other models

There were just a few approaches to simultaneous modelling of the wavelength dependencies of interstellar extinction and linear polarization. In most papers where an imperfect alignment of grains was involved (along with a power-law size distribution) the authors derived that particles with  $r_V \lesssim 0.1 \mu\text{m}$  should be poorly (randomly) oriented while those with  $r_V \gtrsim 0.1 \mu\text{m}$  should be nearly perfectly aligned (e.g., Kim & Martin 1995; Draine & Fraise 2009). Some physical grounds for such alignment of interstellar

grains come, e.g., from Mathis (1986), Goodman & Whittet (1995), Martin (1995).

As the actual composition of carbonaceous grains in the ISM is still unknown (Draine & Fraisse 2009), in interstellar dust modelling one often considered large (sub-micron) graphite grains instead of amorphous carbon ones. As polarization produced by such graphite spheroids may be to a certain extent peculiar, the carbonaceous grains were often assumed to be either spherical or randomly aligned (e.g., Mathis 1986; Draine & Fraisse 2009).

To compare these approaches with our one, we have made some calculations for *three additional models*:

model A1 – imperfectly DG-aligned silicate and *graphite* spheroids;

model A2 – imperfectly DG-aligned silicate spheroids and *randomly oriented* amorphous carbon spheroids;

model A3 – silicate and amorphous carbon spheroids with *randomly oriented small particle* and perfectly DG-aligned large particle.

The grain size distributions and the population of small graphite particles are not varied, i.e. they are the same as in our basic model described in Sects. 2.1–2.4.

When considering graphite spheroids in the A1 model, we use the standard 2/3–1/3 approximation which gives reasonably accurate results (Draine & Malhotra 1993). The orientation function for amorphous carbon grains in the A2 model is  $f(\beta) = \sin \beta$ . In the A3 model this function is size dependent

$$f(\beta, r_V) = \begin{cases} \sin \beta & \text{for } r_V \leq r_{V,\text{cut}} \\ \delta(\beta) & \text{for } r_V > r_{V,\text{cut}} \end{cases}, \quad (14)$$

where  $\delta(\beta)$  is the Dirac delta function,  $r_{V,\text{cut}}$  a cut-off parameter.

So, we assume that all particles whose volume is smaller than that of a sphere with the radius  $r_{V,\text{cut}}$  are randomly oriented, while other particles have perfect rotation alignment, i.e. their angular moment vectors are parallel to the magnetic field direction. In our calculations we choose  $r_{V,\text{cut}}$  to be 0.11  $\mu\text{m}$  (cf. Draine & Fraisse 2009).

Some results obtained with the models considered are presented in Fig. 2, the fitting procedure used is described in the next Section. We see that all models can give very similar extinction curves that well fit the observational data. Only the A1 model with large graphite particles provides the curve that differs a bit from others. Certainly, the extinction data are not enough to constrain the models.

The model dependencies of polarizing efficiency shown in the right panel of the figure vary in form. In particular, the model with randomly oriented carbonaceous particles produces too small polarization in the visual and IR parts of spectrum. Apparently, in this case a better fit can be reached if one takes a more complex size distribution function or (and) increases the alignment efficiency of silicate grains. However, this complicates the model, therefore, we prefer our model having a small number of the parameters and simple assumptions.

### 3 COMPARISON WITH OBSERVATIONAL DATA

So far, most efforts in modelling of the optics of interstellar dust were directed at fitting of the average Galactic extinction curve (often with some additional constraints) by making use of spherical particles. Interpretation of the curves observed towards individual objects was very seldom, and that of the extinction and polarization curves was done only for a few objects. Il'in (1987) applied the model of infinitely long coated cylinders with the imperfect DG-alignment to the interpretation of the wavelength dependence of interstellar extinction, linear and circular polarization for the star HD 204827 in the visual and IR parts of spectrum. Kim & Martin (1994) used infinitely long cylinders with the perfect DG-alignment ( $\Omega = 90^\circ$ ) to interpret the data for HD 25443 and HD 197770. Infinitely long cylinders with a silicate core and organic refractory mantle were chosen by Li & Greenberg (1998) for the interpretation of the extinction and polarization curves for HD 210121. So, Voshchinnikov & Das (2008) were first who utilized the spheroidal model to consider both the extinction and polarization wavelength dependencies observed for two stars.


#### 3.1 Sample of objects

For application of our model we chose stars seen through a single cloud with the extinction and polarization curves known in a wide wavelength region. Initially, preference was given to five stars (HD 24263, HD 62542, HD 99264, HD 147933 and HD 197770) with measured UV polarization (Anderson et al. 1996). These stars are not peculiar and not very distant. Later, two stars (HD 147165 and HD 179406) for which the polarization data in the visible were obtained by us was added to our sample.

In this paper we discuss extinction and polarization of six stars (HD 24263, HD 99264, HD 147165, HD 147933, HD 179406, and HD 197770). Two objects (HD 62542 and HD 147933) were considered in our previous paper (VD08). Almost all the stars are not very far away from the galactic plane (Table 1) and are in dusty environment. One can assume that these stars are seen through single dust clouds as they have weak rotation angle of linear polarization (Anderson et al. 1996) and narrow symmetric sodium D<sub>1</sub> and D<sub>2</sub> lines (Zubko, Krelowski & Wegner 1996). A few comments on the stars should be made.

*HD 24263* is a double star (the secondary is a B9 star) seen through a large cloud AG6 located between the Local Bubble and Orion–Eridanus Super-bubble at distance  $\sim 150$  pc and observed in Na I D and Ca II lines (Genova & Beckman 2003; Welsh et al. 2005). One also detects the molecules CH, CH<sup>+</sup> and CN in the line of sight (Penprase et al. 1990).

*HD 99264* is located in the region of the Scorpius–Centaurus OB association and the Chameleon–Musca dark clouds (Platais et al. 1998; Corradi, Franko & Knude 2004). Consideration of high-resolution Na I D profiles and *uvby* $\beta$  photometry for over 60 nearby B stars allows Corradi, Franko & Knude (2004) to conclude that interstellar gas and dust in this region are distributed in two extended sheet-like structures. The nearby feature located at the distance  $< 60$  pc provides the colour excess  $E(b - y) \approx$



**Figure 2.** Normalized extinction (left panel) and polarizing efficiency (right panel) for our basic model and three additional models with graphite spheroids (A1); randomly oriented amorphous carbon spheroids (A2); small randomly oriented silicate and amorphous carbon spheroids (A3) (see the text for mode details). Oblate spheroids with  $a/b = 3$  and  $\Omega = 33^\circ$ , for other parameters see the last line in Table 3. Observational data for HD 147933.

0.05 while the second feature at 120–150 pc has  $E(b - y) \approx 0.2$ .

*HD 147165* ( $\sigma$  Sco) is a beta Cephei type pulsating variable and a spectroscopic binary with the secondary being a B1V star (North et al. 2007). It is a member of the Upper Scorpius subgroup within the Sco OB2 association related with reflection nebula in the Ophiuchus cloud complex region (Mamajek 2008).

*HD 147933* is a component of the well-studied binary star  $\rho$  Oph AB. High-resolution KI, NaI, CaII observations show very complex profiles of atomic lines in its spectrum (Snow et al. 2008). The star is associated with a strong reflection nebula and a significant amount of dust (and gas) is assumed to lie behind it.

*HD 179406* (20 Aql) is a variable star behind a translucent cloud (Hanson et al. 1992; Dobashi et al. 2005). Though the absorption and emission lines have at least 3 components, the dominant one is seen in most of the atomic and all molecular lines.

*HD 197770* is an evolved spectroscopic eclipsing binary star with both components being B2III stars (Clayton 1996). The object is close to Cyg OB7 and Cep OB2 and probably lies on the edge of a star formation region including molecular clouds L1036 and L1049. It is associated with non-stellar IRAS sources (see for more details Gordon et al. 1998).

So, the clouds observed in the selected lines of sight look to be rather different and hence their dust grain ensemble parameters may also differ.

### 3.2 Observational data

Basic information on the selected stars is presented in Table 1. The distance  $D$ , colour excess  $E(B - V)$ , ratio of total extinction to the selective one  $R_V$  and extinction  $A_V$  for all stars except HD 24263 and 197770 are from Fitzpatrick & Massa (2007), with the data for these two

stars being from Valencic, Clayton & Gordon (2004). The MK spectral types were taken from the catalogue of stellar spectral classifications (Skiff 2009).

The photometric data in the standard visual – near infrared bands were found in the general catalogue of photometric data (Mermilliod et al. 1997) and recent papers referred in the Simbad database (see Table 2). When the JHK data were unavailable, the JHK<sub>S</sub> values from the 2MASS catalogue (Cutri et al. 2003) were used. When the RI data were absent, we applied with caution the data from the USNO B1 catalogue (Monet et al. 2003). To derive the interstellar extinction in the line of sight the intrinsic colours from Straizys (1992) were utilized. The UV extinction parameters were taken from Valencic, Clayton & Gordon (2004) and Fitzpatrick & Massa (2007).

Uncertainties of UV extinction are mainly a result of errors in  $R_V$  and  $A_V$  values. Note that typical relative errors of these quantities are 10–15% (see, e.g., Valencic, Clayton & Gordon 2004). An additional source of relatively large errors of extinction values in the visual and near-IR regions is uncertainty of the intrinsic colors (cf., e.g., Bessell & Brett 1988; Straizys 1992; Ducati et al. 2001). So, relative errors of extinction for individual objects are rather large (see, e.g., Fig. 2).

Table 2 also contains references to the papers with polarimetric data. Additional polarimetric observations of HD 147165 and HD 179406 were performed by one of the authors (HKD) in 2007 at a 2m telescope of the Girawali Observatory (Pune, India). The IUCAA Faint Object Spectrograph and Camera (IFOSC) was used in the polarimetric mode. The polarization parameters were measured in 10 bands in the wavelength region about 0.37 – 0.60  $\mu\text{m}$  and a standard package was applied for data analysis.

Note that for two stars (HD 62542 and HD 197770), far-UV extinction curves were derived from IUE and FUSE observations by Sofia et al. (2005). For two stars (HD 147933 and HD 197770), the important polarization data in the

**Table 1.** Target stars.

Star	$l$	$b$	$D$ (pc)	Sp	$R_V$	$A_V$
HD 24263	182.1	-34.9	171	B5V	3.44	0.72
HD 62542	255.9	-9.2	396	B5V	2.74	0.99
HD 99264	296.3	-10.5	266	B2.5V	3.15	0.85
HD 147165	351.3	+17.0	137	B1III	3.60	1.48
HD 147933	353.7	+17.7	118	B2.5V	4.41	2.07
HD 179406	28.2	-8.3	227	B3V	2.88	0.89
HD 197770	93.9	+9.0	943	B2V	2.77	1.61

**Table 2.** Photometric and polarimetric data sources.

Star	Photometry <sup>a</sup>	Polarimetry <sup>b</sup>
HD 24263	V04 (UV); M97 (UBV); M03 (RI); L05 (JHKL)	A96 (UV & vis-nearIR)
HD 62542	S05 (farUV); F07 (UV); M97 (BV); M03 (RI); C03 (JHK <sub>S</sub> )	A96 (UV & vis-nearIR)
HD 99264	F07 (UV); M97 (UBVRI); C03 (JHK <sub>S</sub> )	A96 (UV & vis); S69 (UBV)
HD 147165	V04 (UV); W02 (UBVRIJHKLM)	this work (vis); C66 (vis-nearIR)
HD 147933	F07 (UV); M97 (UBVRI); C03 (JHK <sub>S</sub> )	A96 (UV-nearIR)
HD 179406	V04 (UV); W02 (UBVJHKL); M03 (RI)	this work (vis)
HD 197770	S05 (farUV); V04 (UV); M97 (UBV); M03 (RI); C03 (JHK <sub>S</sub> )	A96 (UV-nearIR)

<sup>a</sup> M97 is for Mermilliod et al. (1997), W02 for Wegner (2002), C03 for Cutri et al. (2003), M03 for Monet et al. (2003), V04 for Valencic, Clayton & Gordon (2004), L05 for Larson & Whittet (2005), S05 for Sofia et al. (2005), F07 for Fitzpatrick & Massa (2007).

<sup>b</sup> C66 is for Coyne & Gehrels (1966), S69 for Serkowski & Robertson (1969), A96 for Anderson et al. (1996).

J, H and K bands are also available (Wilking et al. 1980; Wilking, Lebofsky & Rieki 1982).

### 3.3 Fitting procedure and parameter variations

A general problem of dust optics modelling is in obtaining the absolute values of the model parameters. As it has been noted many times (e.g., Greenberg 1978), similar extinction occurs when a product of the typical particle size  $\langle r \rangle$  and the particle refractive index does not change, i.e.

$$\langle r \rangle |m - 1| \approx \text{const.}$$

The average Galactic extinction curve was successfully reproduced using quite different mixtures of particles with exponential (Hong & Greenberg 1980), power-law (Mathis, Rumpl & Nordsieck 1977) and more complicated grain size distributions (Mathis 1996; Weingartner & Draine 2001). Effects of variations of the distribution parameters (e.g.,  $r_{V,\min}$ ,  $r_{V,\max}$ ,  $q$  for a power-law distribution) on extinction are well studied for spheres (see, for example, Voshchinnikov & Il'in 1993).

Joint modelling of interstellar extinction and polarization adds at least one parameter — particle shape — to the model. In the case of spheroids the particle shape is defined by the spheroid's aspect ratio  $a/b$ . Besides this, we can consider prolate (“cigars”) and oblate (“pancakes”) spheroids. If such particles are imperfectly oriented, the alignment parameters (in our case,  $\delta_0$  and  $\Omega$ ) very weakly affect extinction. Note that the normalized extinction  $A_\lambda/A_V$  essentially changes in the UV region when  $a/b$  varies (see Fig. 3 in VD08).

In order to analyze importance of the model parameters we reproduced the observational data available for HD 147933 with different particles. Some results are shown in Figs. 2 and 3 (see also Fig. 8 in VD08), the parameters of the fits are given in Table 3. Note that in VD08 the fitting was simply “chi-by-eye” while in this paper we apply minimization of  $\chi^2$ .

The fitting approach used included two steps: i) finding of the size distribution parameters and the relative column densities  $\mathcal{K}_i$  from approximation of the extinction data using particles of given shape, and ii) a more accurate determination of the alignment parameters  $\delta_0$  and  $\Omega$  from fitting of the wavelength dependence of polarizing efficiency  $P(\lambda)/A(\lambda)$ .

It is interesting that by varying the parameters  $r_{V,\min}$ ,  $r_{V,\max}$  and  $q$  and the relative density of carbon ( $\mathcal{K}_C$ ) and silicate ( $\mathcal{K}_{Si}$ ) grains one can reproduce the observational data with prolate as well as oblate particles of different aspect ratios. Note that after fixing the particle type (prolate/oblate) and  $a/b$  one can estimate only the values of  $\mathcal{K}_C$ ,  $\mathcal{K}_{Si}$ ,  $r_{V,\min}$  and  $q$  as the extinction and polarization curves is little affected by variation of  $r_{V,\max}$ .


The alignment degree for three models presented in Table 3 is found to be slightly larger than the standard interstellar value ( $\delta_0 \approx 0.2 \mu\text{m}$ , Voshchinnikov 1989). This can be a result of enhanced magnetic fields in the interstellar cloud near  $\rho$  Oph with which HD 147933 is related. The most exciting conclusion of our modelling is a close coincidence of the magnetic field directions — the values of  $\Omega$  for various fits differ less than  $10^\circ$ . We suggest that this model parameter can be more or less reliably determined from such modelling.

Possibly, a better coincidence of the theory with the ob-

**Table 3.** Modelling results for HD 147933.

$\mathcal{K}_C$	amorphous carbon				astrosil			graphite	shape;	$\delta_0(\mu\text{m})$	$\Omega$ (deg)	$\chi^2$	
	$r_{V,\text{min}}^a$	$r_{V,\text{max}}$	$q$	$\mathcal{K}_{\text{Si}}$	$r_{V,\text{min}}$	$r_{V,\text{max}}$	$q$	$\mathcal{K}_{\text{gra}}$	$a/b$			$A/A_V$	$P/A$
0.24	0.03	0.15	1.5	0.73	0.08	0.25	1.5	0.03	pro; 5	0.5	35	1.04	24.6
0.54	0.10	0.20	2.0	0.40	0.01	0.25	1.3	0.06	pro; 3	0.3	42	0.19	8.2
0.50	0.10	0.25	2.0	0.47	0.07	0.20	2.0	0.03	obl; 3	0.3	33	0.19	2.2

<sup>a</sup>  $r_{V,\min}$  and  $r_{V,\max}$  are in  $\mu\text{m}$



**Figure 3.** Comparison of the normalized extinction (left panel) and the polarizing efficiency (right panel) observed for HD 147933 with results of modelling. The solid curves show the case of prolate ( $a/b = 3$ ) spheroids having a power-law size distribution. Other model parameters are given in Table 3.

servations could be obtained if we consider different alignment parameters for silicate and the carbonaceous grains. Certainly, consideration of non-aligned carbonaceous grains cannot be ruled out (Li & Greenberg 2002; Chiar et al. 2005; see, however, Fig. 2).

### 3.4 Fitting results and discussion

The normalized extinction  $A(\lambda)/A_V$  and the polarizing efficiency  $P(\lambda)/A(\lambda)$  were calculated with the theory of Sect. 2 and compared with the observational data available for the stars mentioned in Sect. 3.1. The results of this comparison are plotted in Figs. 3–4 (see also Fig. 9 in VD08 for HD 62542) with the parameters of the fits being presented in Table 4. We start with comments on the individual objects and then make general conclusions.

*HD 62542.* The UV extinction curve has a very broad and weak 2175 Å bump and a steep far-UV rise (Snow & McCall 2006). Our model suggests that the major part of grains are small carbon particles with a very narrow size distribution.


*HD 99264.* The extinction curve for it is characterized by  $R_V$  of about 3 (Fitzpatrick & Massa 2007), the UV extinction curve is a bit “shallow” but not anomalous (Carnochan 1986). The polarization in an UV region and the UBV bands is well described by Serkowski curve with  $\lambda_{\max} = 0.55\mu\text{m}$  (Anderson et al. 1996; Martin, Clayton & Wolff 1999). So,

we have the case close to the average Galactic curves, but certainly there are not enough polarization data for sound comparison.

*HD 147165* and *HD 179406.* Extinction in these lines of sight along with element abundances were interpreted by Zubko, Krelowski & Wegner (1996, 1998) using rather wide size distributions of grains. These authors applied spherical particles of different materials as well as inhomogeneous particles, which makes impossible a direct comparison of their results with those obtained in our work.

*HD 197770.* For this star the polarizing efficiency is maximal: it reaches  $P/A = 3.7$  at the J band ( $\lambda = 1.25\mu\text{m}$ ). The polarization is enhanced in the region of the UV bump (super-Serkowski behaviour) but strong UV bump in extinction results in no structure for  $P/A$  curve (see insert in Fig. 4). Our model shows the largest fraction of graphite in the grain mixture for this star.

Thus, we have fitted the observational data using prolate or oblate spheroids with  $a/b = 2$  or 3 and intermediate values of the alignment parameter  $\delta_0 = 0.3 - 0.5\mu\text{m}$ . Contributions of carbonaceous and silicate grains to extinction and polarization are nearly equal excluding the case of HD 62542 for which extinction is very peculiar. A contribution of small graphite particles to extinction usually is less than 10%. The parameters of the size distributions for silicate and amorphous carbon grains differ from the standard



**Figure 4.** Comparison of observed and modelled normalized extinction (left panel) and polarizing efficiency (right panel) for five stars. The solid curves show results of calculations. The model parameters are given in Table 4.



**Table 4.** Modelling results.

Star	$\mathcal{K}_C$	amorphous carbon				astrosil			graphite	shape;	$\delta_0(\mu\text{m})$	$\Omega$ (deg)
		$r_{V,\text{min}}^a$	$r_{V,\text{max}}$	$q$	$\mathcal{K}_{\text{Si}}$	$r_{V,\text{min}}$	$r_{V,\text{max}}$	$q$	$\mathcal{K}_{\text{gra}}$	$a/b$		
HD 24263	0.48	0.08	0.40	3.5	0.47	0.04	0.15	2.1	0.05	pro; 2	0.3	60
HD 62542	0.85	0.07	0.10	2.7	0.10	0.005	0.25	2.7	0.05	pro; 2	0.5	40
HD 99264	0.44	0.09	0.30	6.0	0.50	0.04	0.18	2.0	0.06	pro; 3	0.5	50
HD 147165	0.49	0.06	0.30	2.2	0.44	0.04	0.20	2.2	0.07	pro; 2	0.5	54
HD 147933	0.50	0.10	0.25	2.0	0.47	0.07	0.20	2.0	0.03	obl; 3	0.3	33
HD 179406	0.45	0.10	0.40	6.0	0.47	0.04	0.18	2.5	0.08	pro; 3	0.5	49
HD 197770	0.43	0.05	0.20	2.8	0.46	0.01	0.17	1.3	0.11	pro; 3	0.5	42

<sup>a</sup>  $r_{V,min}$  and  $r_{V,max}$  are in  $\mu m$

MRN mixture (see Sect. 2.2). As usual, very small particles with  $r_{V,min} \lesssim 0.01 \mu m$  are absent in the grain ensembles. Besides this, in many cases the slope of the  $n(r_V)$  dependence is less steep than that for the MRN mixture (i.e.  $q < 3.5$ ).

The inclination of the magnetic field direction relative to the line of sight determined for all target stars lies in the interval  $\Omega \approx 30 - 60^\circ$ . These values of  $\Omega$  are derived rather accurately because of a strong dependence of  $P/A$  on  $\Omega$  (see Figs. 4 and 6 in VD08). The magnetic fields seem to be connected with local dust clouds in the directions to the selected stars. At the same time, the obtained values of  $\Omega$  do not contradict the general pattern of the galactic magnetic field in the solar neighbourhood as found from pulsar measurements (Vallée 2004, 2008).

It should be noted that one usually assumed that small grains were badly oriented as the interstellar polarization degree decreases with a growing  $\lambda^{-1}$  in the UV region (Whittet 2003). However, for all the objects we have fitted the observational data by using the simple size and orientation distributions with small particles being rather well aligned. To see that one should consider Eqs. (11)–(13) and keep in mind that  $\delta_0$  is about 0.4 and the temperature ratio about 0.1. For  $r_V \sim 0.06 \mu m$ , this gives  $\xi \sim 0.5$  when  $\xi = 0$  corresponds to the perfect alignment while  $\xi = 1$  means the random orientation.

## 4 CONCLUSIONS

We have extended the spheroidal model of interstellar dust by considering imperfectly dynamically aligned amorphous carbon and silicate prolate and oblate spheroids with a power-law size distribution and a small fraction of graphite spheres involved to explain the 2175Å bump. Our model being as simple as possible was applied to fit the normalized extinction  $A(\lambda)/A_V$  and the polarizing efficiency  $P(\lambda)/A(\lambda)$  observed in the near IR to far UV region.

In contrast to earlier works we exactly calculated the optical properties of spheroids, for the first time utilized their imperfect Davis–Greenstein alignment and applied the spheroidal model to fit the data obtained for individual stars instead of the commonly used average Galactic curves.

A comparison of our model with those recently suggested by other authors has demonstrated that the extinction data are well fitted by all the models while only our model allows one to satisfactorily fit the polarization data

with simple size and orientation distributions. Contrary to other works we find that the data can be well fitted with aligned carbonaceous and small silicate non-spherical grains.

Investigation of our model has shown that the only parameter that can be determined more or less accurately from observational data fitting is the direction of magnetic field relative to the line of sight. As the projection of the magnetic field on the sky plane is given by the positional angle of polarization  $\vartheta$ , multi-wavelength extinction and polarization observations can be used for study of the spatial structure of interstellar magnetic fields.

Variations of the interstellar extinction and polarization curves for the considered stars can be interpreted by differences in the relative fractions of carbon and silicate grains and their size distributions in the clouds crossed by the lines of sight with the alignment parameter having intermediate values  $\delta_0 = 0.3 - 0.5 \mu m$  (for the particle aspect ratio  $a/b = 2 - 3$ ). We assume that ambiguous results could be ruled out by considering dust phase abundances.

## ACKNOWLEDGMENTS

One of the authors (HKD) is deeply indebted to the IUCAA's instrumentation and observatory team for their support and cooperation extended during the course of this work. The work was partly supported by the grants RFBR 07-02-00831, NTP 2.1.1/665 and NSh 1318.2008.2.

## REFERENCES

- Aitken D.K., 1996, ASP Conf. Ser., 97, 225
- Anderson C.M., Weitenback A.J., et al., 1996, AJ, 112, 2726
- Bessell M.S., Brett J.M., 1988, PASP, 100, 1134
- Carnochan D.J., 1986, MNRAS, 219, 903
- Chiar J. E., Adamson A. J., Kerr T. H., Mason R., et al., 2005, ASP Conf. Ser., 343, 352
- Clayton G.C., 1996, PASP, 108, 401
- Corradi W.J.B., Franco G.A.P., Knude J., 2004, MNRAS, 347, 1065
- Coyne G.V., Gehrels T., 1966, AJ, 71, 355
- Cutri R.M., Skrutskie M.F., Van Dyk S., Beichman C.A., et al., 2003, 2MASS All-Sky Catalog of Point Sources, Univ. Massach. & IPAC / CIT
- Das H.S., Sen A.K., 2006, A&A, 459, 271




Figure 4. (continued).

Davis L., Greenstein J.L., 1951, *ApJ*, 114, 206  
 Dobashi K., Uehara H., Kandori R., Sakurai T., Kaiden M., Umemoto T., Sato F., 2005, *PASJ*, 57, S1  
 Draine B.T., 2003, *ARA&A*, 41, 241  
 Draine B.T., 2009, *ASP Conf. Ser.*, in press ([arXiv:0903.1658])  
 Draine B.T., Allaf-Akbari K., 2006, *ApJ*, 652, 1318  
 Draine B.T., Fraisse A.A., 2009, *ApJ*, 696, 1  
 Draine B.T., Lee H.M., 1984, *ApJ*, 285, 89  
 Draine B.T., Malhotra S., 1993, *ApJ*, 414, 632  
 Ducati J.R., Bevilacqua C.M., Rembold S.B., Ribeiro D., 2001, *ApJ*, 558, 309  
 Fitzpatrick E.L., Massa D.L., 2007, *ApJ*, 663, 320  
 Genova R., Beckman J.E., 2003, *ApJS*, 145, 355  
 Goodman A.A., Whittet D.C.B., 1995, *ApJ*, 455, L181  
 Gordon K.D., Clayton G.C., Smith T.L., et al., 1998, *AJ*, 115, 2665  
 Greenberg J.M., 1978, *Cosmic Dust*, ed. by J.A.M. McDonnell, p. 187  
 Greenberg J.M., Li A., 1996, *New extragalactic perspec-*

*tives in the New South Africa*, ed. D.L. Block and J.M. Greenberg, Kluwer, Dordrecht, p. 118  
 Gupta R., Mukai T., Vaidya D.B., Sen A.K., Okada Y., 2005, *A&A*, 441, 555  
 Hanson M.M., Snow T.P., Black J.H., 1992, *ApJ*, 392, 571  
 Hildebrand R.H., Dragovan M., 1995, *ApJ*, 450, 663  
 Hong S.S., Greenberg J.M., 1980, *A&A*, 88, 194  
 Iati M.A., Saija R., Borghese F., Denti P., Cecchi-Pestellini C., Williams D.A., 2008, *MNRAS*, 384, 591  
 Il'in A.E., 1987, *Afz*, 27, 477  
 Jones A.P., 2000, *J. Geophys. Res.*, 105, 10257  
 Kim S.H., Martin P.G., 1994, *ApJ*, 431, 783  
 Kim S.H., Martin P.G., 1995, *ApJ*, 444, 293  
 Krełowski J., 1989, *Astron. Nachr.*, 310, 281  
 Laor A., Draine B.T., 1993, *ApJ*, 402, 441  
 Larson K.A., Whittet D.C.B., 2005, *AJ*, 623, 897  
 Lasue J., Levasseur-Regourd A.C., Fray N., Cottin H., 2007, *A&A*, 473, 641  
 Lazarian A., 2007, *J. Quant. Spectr. Rad. Transf.*, 106, 225  
 Lee H.M., Draine B.T., 1985, *ApJ*, 290, 211

- Li A., Greenberg J.M., 1997, *A&A*, 323, 566
- Li A., Greenberg J.M., 1998, *A&A*, 339, 591
- Li A., Greenberg J.M., 2002, *ApJ*, 577, 789
- Li A., Chen J.H., Li M.P., et al., 2008, *MNRAS*, 390, L39
- Li M.P., Zhao G., Li A., 2007, *MNRAS*, 382, L26
- Mamajek E.E., 2008, *Astron. Nachr.*, 329, 10
- Martin P.G., 1978, *Cosmic Dust*, Oxford Univ. Press, Oxford
- Martin P.G., 1995, *ApJ*, 445, L63
- Martin P.G., Clayton G.C., Wolff M.J., 1999, *ApJ*, 510, 905
- Mathis J.S., 1986, *ApJ*, 308, 281
- Mathis J.S., 1996, *ApJ*, 472, 643
- Mathis J.S., Rimpl W., Nordsieck K.H., 1977, *ApJ*, 217, 425
- Mermilliod J.-C., Mermilliod M., Hauck B., 1997, *A&AS*, 124, 349 (<http://obswww.unige.ch/gcpd/gcpd.html>)
- Min M., Hovenier J.W., Dominik C., et al., 2006, *J. Quant. Spectr. Rad. Transf.*, 97, 161
- Monet D.G., Levine S.E., Canzian B., Ables H.D., et al., 2003, *AJ*, 125, 984
- Moreno F., Munoz O., et al., 2007, *J. Quant. Spectr. Rad. Transf.*, 106, 348
- North J.R., Davis J., Tuthill P.G., et al., 2007, *MNRAS*, 380, 1276
- Onaka T., 1980, *Ann. Tokyo Astron. Obs.*, 18, 1
- Penprase B.E., Blades J.C., Danks A.C., Crane P., 1990, *ApJ*, 365, 241
- Platais I., Kozhurina-Platais V., van Leeuwen F., 1998, *AJ*, 116, 2423
- Rogers C., Martin P.G., 1979, *ApJ*, 228, 450
- Rouleau F., Martin P.G., 1991, *ApJ*, 377, 526
- Serkowski K., Robertson J.W., 1969, *ApJ*, 158, 441
- Skiff B.A., 2009, *Catalogue of Stellar Spectral Classifications*, Lowell Obs.
- Snow T.P., McCall B.J., 2006, *Ann. Rev. Astron. Astrophys.*, 44, 367
- Snow T.P., Destree J.D., Welty D.E., 2006, *ApJ*, 679, 512
- Sofia U.J., Wolff M.J., Rachford B., Gordon K.D., et al., 2005, *ApJ*, 625, 167
- Stark C.R., Potts H.E., Diver D.A., 2006, *A&A*, 457, 365
- Straizys V., 1992, *Multicolor Stellar Photometry*, Pachart Publ. House, Tucson
- Vaidya D.B., Gupta R., Dobbie J.S., Chylek P., 2001, *A&A*, 375, 584
- Vaidya D.B., Gupta R., Snow T.P., 2007, *MNRAS*, 379, 791
- Vallée J.P., 2004, *New Astron. Rev.*, 48, 763
- Vallée J.P., 2008, *ApJ*, 681, 303
- Valencic L.A., Clayton G.C., Gordon K.D., 2004, *ApJ*, 616, 912
- Voshchinnikov N.V., 1989, *Astron. Nachr.*, 310, 265
- Voshchinnikov N.V., 2004, *Astrophys. Space Phys. Rev.*, 12, 1
- Voshchinnikov N.V., Das H.K., 2008, *J. Quant. Spectr. Rad. Transf.*, 109, 1527 (VD08)
- Voshchinnikov N.V., Farafonov V.G., 1993, *Astrophys. Sp. Sci.*, 204, 19
- Voshchinnikov N.V., Farafonov V.G., 2003, *J. Comp. Math. Math. Phys.*, 43, 1299
- Voshchinnikov N.V., Il'in V.B., 1993, *Sov. Astron.*, 37, 21
- Voshchinnikov N.V., Il'in V.B., Henning Th., Dubkova D.N., 2006, *A&A*, 445, 167
- Wegner W., 2002, *Balt. Astron.*, 11, 1
- Weingartner J.C., Draine B.T., 2001, *ApJ*, 548, 296
- Welsh B.Y., Sallmen S., Jelinsky S., 2005, *A&A*, 440, 547
- Whittet D.C.B., 2003, *Dust in the galactic environment*, Bristol:IOP
- Willing B.A., Lebofsky M.J., et al., 1980, *ApJ*, 235, 905
- Willing B.A., Lebofsky M.J., Rieke G.H., 1982, *AJ*, 87, 695
- Wolf S., Voshchinnikov N.V., Henning Th., 2002, *A&A*, 385, 365
- Zubko V.G., Krelowski J., Wegner W., 1996, *MNRAS*, 283, 577
- Zubko V.G., Krelowski J., Wegner W., 1998, *MNRAS*, 294, 548